




The Potassium–Argon Laser Experiment (KArLE): Design Concepts Yuichiro Cho^{1,2} & B.A. Cohen³ NASA Marshall Space Flight Center, yuichiro.cho@nasa.gov, ²Univ. Alabama in Huntsville, ²NASA Goddard Space Flight Center



1. Background

♦When were they formed?




▲ Beautiful stratigraphy observed by Curiosity on Mars. Nobody knows exactly when these layers were formed. Knowing there absolute ages is important to understand the processes that emplaced these



▲An important goal of the science community is to measure the age of various key geologic units to reconstruct Martian history and correlate it with other Solar System events.

2. In situ K-Ar dating



▲ Schematics of KArLE measurements. Spot-by-spot analyses yield the isochron age of target rocks, enhancing the reliability and accuracy of the K-Ar age measurement.

K-Ar dating with LIBS-MS approach

(Cho+ 2016, PSS: Cohen+ 2014, GGR: Devismes+ 2016, GGR)

$$\text{K-Ar age } t = \frac{1}{\lambda} \ln \Bigl(\frac{\lambda}{\lambda_{\rm o}} \frac{[^{40} {\rm Ar}]_{\rm rad}}{[^{40} {\rm K}]} + 1 \Bigr)$$

- 1. Laser ablates a target rock in vacuum chamber
- 2. K contents measured with LIBS (e.g., ChemCam)
- 3. Released Ar measured using mass spectrometry (e.g., SAM)
- 4. K and Ar related by volume of the ablated pit using optical measurement (e.g., MAHLI)
- Use TRL 9 components to achieve new science
- 1. Payload synergy
- 2. Reasonable cost
- 3 Low risk
- 4. Near-term implementation

3. KArLE breadboards



- ▲ KArLE breadboard at MSFC
- HR2500+ Ocean Optics spectrometer
- slumn for a camera recording the adeview of the plasma
- 5- Vacuum line including getter, preumatic values, furbonicleoular
- ▲ Breadboard in Japan / Field test at Izu island
 - 1- MausPro 2000 Ocean Optics spectrometer
- 3. KOORE AO. 1010 lane
- Ablation cell with sample handle
- Vecoum line including getter, manual valves, B.A.gauge 9- Mains Spectrometer (America GMS)
- Mass Spectrometer (Hiden Analytics QMS.) hat Detect (TMS)


4. KArLE flight concepts (examples)

KArLE enables various configurations, because

- KArLE is a partner-provided instrument suite; agnostic to specific analysis providers
- KArLE allows for flexible implementation with multiple sample delivery systems (e.g., core, pebble, and slab)
- KArLE-specific hardware is mechanically simple sample handling system (SHS). which provides vacuum sealing and laser pits observation.



Curiosity-like configuration. The mast mounted LIBS measures the sample in the KArLE chamber. The chamber could be mounted on the deck or inside the rover body. The gas processing system and the mass spectrometer is stored in the rover body.



Point design for candidate KArLE flight configuration.

- LIBS+MS+camera payload -15 kg; distributed volume;
- power = 10 65 W Chamber+sample handling -variable but probably -1-2 kg: low power
- (stepper motors, gasket preload)

◆The amount of

residual 47Ar

when different

elastomeric Orings are used

under various


Viton Orings

can maintain

low 40 Ar blank

ambient


conditions








▲ KArLE Sample Handling System (SHS). The SHS must be capable of ingesting a sample. achieving a vacuum seal, and enabling the measurements to be performed on the enclosed sample. A SAM-like elevator actuator seals the chamber. Samples are introduced into and ejected from the chamber by a spoon-like manipulator.




▲An alternative approach: Compact box configuration. Entire geochronology package is installed in the rover/lander body [Cho+ 2016b, Trans, JSASS]. Our experiments suggest that elastomeric O-rings would be useful for Ar analyses (Cho+ 2017, submitted to ASR).

5. Results of LIBS experiments




- (a) Emission spectrum of a standard sample
- (b) Calibration curve at the laser pulse energy of 30 mJ. Best-fit calibration curve and 10 prediction bands are shown.
- (c) Errors in K-Ar age measurement as a function of K contents. Error propagation was calculated assuming 10% error in 40Ar measurements.





- Measurement error was <20% when K₂O >2400 ppm.
- For 30 mJ, normalizing I_{KNorm} with I_{CITTINE} gave the best
- For 15 mJ, normalizing I_{KTRIPPS} with I_{continue} gave the best results
- · With a 30-mJ laser, K-Ar age can be measured with an error of 8% for a 4 Ga rock containing 3000 ppm K₂O.

6. Obtained Isochrons










- ▲K-Ar isochron for the hornblende-biotite gneiss. The data points follow one regression line well, suggesting the viability of isochron measurements with the LIBS-MS approach. The slope and intercept yielded a K-Ar age of 700±190 Ma and an initial ** Ar/26 Ar ratio of 440±280,
- ◆ ** Ar-K plot for (a) homblende-biotite gneiss and (b) pyroxene gneiss samples. The "isochron" slopes agree with the K-Ar ages determined for biotite separates with conventional methods

7. Performance of K-Ar dating

- ►Compiled K-Ar dating results published from multiple labs. Results from multiple laboratories yield whole-rock ages within error of accepted ages and precision close to theoretical
- = TR1, 4 (validation in the laboratory)

Data navner: Sold 2014 Champai Geol; Conen x 2014 Davidanitiests and Sectanalytical Phosper KOprismay + 2018 Sectionships and Geograp/Intel Respect, Chic + 2016 IRSS



8. Work in progress

- Using the laboratory breadboard to measure Mars and Moon analog materials.
- Characterizing and optimizing the performance of the components
- · Pursuing funding for construction and test of the flight concepts

- AND READ AND COLD FOR SOME COLD AND COL